

(11) EP 0 837 074 A2

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
22.04.1998 Bulletin 1998/17

(51) Int. Cl.⁶: C07K 14/78

(21) Application number: 97116293.8

(22) Date of filing: 18.09.1997

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV RO SI

(30) Priority: 19.09.1996 JP 248247/96

(71) Applicant:
HISAMITSU PHARMACEUTICAL CO., INC.
Tosu-shi Saga-ken (JP)

(72) Inventors:
• Fukai, Fumio
Tokorozawa-shi, Saitama 359 (JP)

• Katayama, Takashi
Tokyo 169 (JP)

(74) Representative:
Grünecker, Kinkeldey,
Stockmair & Schwanhäusser
Anwaltssozietät
Maximilianstrasse 58
80538 München (DE)

Remarks:


The applicant has subsequently filed a sequence
listing and declared, that it includes no new matter.

(54) **Biologically active fibronectin fragment as cancer metastasis inhibitor**

(57) The biologically active peptide of the invention
has a number of amino acids of 30 or less, comprises
an amino acid sequence described as sequence ID No.
13 (Tyr Thr Ile Tyr Val Ile Ala Leu) in the sequence listing

and has cell adhesion inhibition activity. The peptide of
the invention has short length and, therefore, synthesis
and handling of the peptide are easy.

Fig.2



EP 0 837 074 A2

Descripti n

FIELD OF THE INVENTION

5 This invention relates to a biological active peptide having cell adhesion inhibition activity.

BACKGROUND OF THE INVENTION

Many attempts have been done to establish a therapy for cancer, and such attempts have shown some advanta-
10 geous therapeutic effects on several cancers. However, a method preventing cancer metastasis has not been estab-
lished as yet.

Surgical therapy, radiotherapy and chemotherapy can be listed as a major method for therapy when one is diag-
nosed as getting cancer. When a patient shows clear abnormal symptom, most cancer probably becomes progressive
and it is difficult to cure it. We can strongly believe that such difficulties is due to cancer metastasis and growing. Con-
15 sequently, patient's condition after being cured mainly depends on the degree of cancer metastasis (Welch et al., Intern.
J. Cancer, 43, 449, 1989). It is extremely difficult to cure cancer completely so long as such a cancer metastasis and
growing are not suppressed. For example, in the case that stomach cancer becomes progressive, it is often that the
cancer has already metastasized to lymph node, liver, lung and so forth. Such a phenomenon is occurred in breast can-
cer, it is reported that progressive cancer has metastasized to lymph node, which is adjacent to it, as well as bone mar-
20 row and lungs. Particularly, in regard to the epithelial cancer, the patient's condition after being cured depends on the
degree of metastasis of cancer to lymph node (Shigeru Tsukakoshi, chemistry to cure cancer, Japan pharmaceutical
society Pharmacia Review, No. 6).

Due to the above reasons, many attempts to elucidate detailed mechanism of cancer metastasis and apply its
mechanism to therapy have done.

25 Cancer metastasis is constituted a complicated process in which cancerous cells departure from primary spot, infil-
trate to peripheral tissues and grow in that metastasized tissues. This process is influenced by both factors of cancerous
cell and host cell. In the former, the primary spot suffers from continuous gene alteration and consists of an uniformity
cell mass showing various cytomorphosis, such as proliferation potency, drug sensitiveness, immunogenicity, configu-
ration, etc. As factors of cancerous cell, metastasis ability, infiltration ability to periphery tissues, production of self
30 growth factors from metastatic cancerous cell, and the like influence cancer metastasis. As factors of host cell, physical
and anatomical condition in the blood, defense mechanism of immunocompetent cell, adhesion interaction between
endothelial cell and blood platelet, growth factors from periphery metastasized tissues, and the like influence cancer
metastasis.

It seems that on the serial process for cancer metastasis, expression and regulation of metastatic properties of can-
cerous cell is influenced by the interaction between cancerous cell and normal host cell, the interaction of cancerous
35 cells with each together, and the interaction between cancerous cell and various biological components such as cell
external matrix. Adhesion between cancerous cells with each together or between cancerous cell and normal cell is
mediated by a cell adhesion molecule expressed on cell surface, and, as such a cell adhesion molecule, cadherin
molecular group (Masatoshi Takeichi, molecular basis for tissue construction - role of cadherin for cell recognition - bio-
chemistry, 59, 1, 1987), immune globulin molecular group, secretin molecular group (Springer, T. A., Nature, 346, 425,
40 1990) and so forth are known. On the other hand, adhesion molecule on cell surface including integrin molecular group
contributes adhesion between cancerous cell and external matrix including glycoprotein such as fibronectin, laminin
and collagen, and proteoglycan being bound with heparan sulfate, chondroitin sulfate etc.

Recently, it was made clear that cell external matrix molecule such as fibronectin or laminin strongly participated in
45 cancer metastasis. It is shown that, after cell suffers from malignant transformation by oncogenic virus or chemical car-
cinogenesis agent etc., synthesis or accumulation into cell external of cell external matrix molecule, especially fibronec-
tin, is decreased (Yamada, K. M. and Olden, K., Nature, 275, 179, 1978), expression of integrin molecule is decreased,
and its adhesion ability to fibronectin is decreased (Plantefaber, L. C. and Hynes, R. O., Cell, 56, 281, 1989). It is also
reported that fibronectin and laminin promote adhesion and metakinesis ability of cancerous cell (McCarthy, J. B., J.
50 Cell Biol., 98, 1474, 1984).

Fibronectin and laminin molecule have domain configuration and let lots of functions disperse in the molecule. Con-
figuration is made clear by biochemical or genetic engineering procedure, and existence of cell adhesion portion and
receptor for it are discovered. RGD sequence in cell binding domain of fibronectin is cell recognition sequence (Piersch-
bacher, M. D., Nature, 309, 30, 1984), and the sequence exists and functions in lots of cell adhesion related molecules,
55 and the receptor for the sequence is identified as alpha 5 beta 1 integrin receptor (Pytela, R., Cell, 40, 191, 1985).

Adhesion interaction between cell adhesion molecule and cell has been made clear, and recently application of cell
adhesion peptide to inhibit cancer metastasis has been attempted. Humphries et al. show that GRGDS peptide as an
adhesion signal of fibronectin inhibits experimental metastasis of melanoma cell to lung (Humphries, M. J., Science,

233, 467, 1986).

However, there are lots of problems to be solved in which clearance rate of the peptide in the blood after administration is fast and the peptide tends to be easily digested by enzyme for applying the peptide as a medicine (Saiki, I. et al., Jpn. J. Cancer Res., 84, 558, 1993).

Exploration of cancer metastasis inhibition materials as mentioned above is necessary to cure cancer completely. However, no substance being excellent medicine for cancer has obtained up to now.

SUMMARY OF THE INVENTION

In the light of the foregoing demand, the present invention was made. The purpose of the present invention is to provide a new peptide having cell adhesion inhibition activity.

The inventors have made an extensive study to resolve the above problems, and found a new peptide having cell adhesion inhibition activity, especially a peptide including the amino acids sequence described in the sequence ID No. 13 of the sequence listing.

The present invention is a peptide which has the number of amino acids of 30 or less, comprises the amino acid sequence described as the sequence ID No. 13 in the sequence listing and has cell adhesion inhibition activity.

The present invention is also a peptide which comprises the amino acid sequence described as the sequence ID No. 13 in the sequence listing and has cell adhesion inhibition activity and which is characterized by having the number of amino acids of 13 - 30.

Further, the present invention is a peptide which has the number of amino acids of 30 or less, comprises the amino acid sequence described as the sequence ID No. 8 in the sequence listing and has cell adhesion inhibition activity.

Furthermore, the present invention is a peptide which has the number of amino acids of 30 or less, comprises the amino acid sequence described as the sequence ID No. 11 in the sequence listing and has cell adhesion inhibition activity.

Still further, the present invention is a peptide which has the number of amino acids of 30 or less, comprises the amino acid sequence described as the sequence ID No. 12 in the sequence listing and has cell adhesion inhibition activity.

The present invention is also a peptide derived from the above-mentioned peptides by insertion, deletion or substitution of one or several amino acids and has the cell adhesion inhibition activity.

In addition, the present invention is a cancer metastasis inhibitor comprising at least one peptide of the above-mentioned peptides.

The present invention will be more fully understood from the detailed description given hereinbelow and the accompanying drawings, which are given by way of illustration only and are not to be considered as limiting the present invention.

Further scope of applicability of the present invention will become apparent from the detailed description given hereinafter. However, it should be understood that the detailed description and specific examples, while indicating preferred embodiments of the present invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the present invention will be apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the sequence of Hep 2 fragment and a position thereon of each peptide used in Experiment 1. The numeral of amino acid shows its position on fibronectin.

Fig. 2 shows the relationship among each peptide used in Experiment 2.

Fig. 3 shows the result of the evaluation of cell adhesion inhibition activity of each KLH modified synthetic peptide.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is a new peptide having cell adhesion inhibition activity, and is characterized by including the amino acids sequence described in the sequence ID No. 13 of the sequence listing. The peptide of the present invention is very short length peptide and, therefore, the peptide can be synthesized easily and provides strong cell adhesion inhibition activity without any denaturation treatment such as urea treatment.

The amino acids sequence of the peptide of the present invention, i.e. amino acids sequence described in the sequence ID No. 13 of the sequence listing, is the same as the amino acids sequence of 1848-1855 portion of fibronectin.

It is reported that fibronectin has cell adhesion activity (Hynes, R. O. et al., J. Cell Biol., 95, 369-377, 1982). On the other hand, it is reported that 30 kDa fragment (Hep 2) derived from Hep 2 domain of fibronectin becomes to have

strong cell adhesion inhibition activity after being denaturated by urea (Fukai, F. et al., Biochemica. Biophys. Res. Commun., 220, 394, 1996). However, it did not suggest or indicate that which amino acid sequence contributes the cell adhesion inhibition activity of Hep 2. Furthermore, Hep 2 consists of 271 amino acids. As a result, the attempt in which Hep 2 itself will be used as a medicine is not desirable because Hep 2 has a high molecular weight and includes non related sequence of cell adhesion inhibition activity. The synthesis and preparation of Hep 2 is difficult because of its high molecular weight, and it is likely that an amino acid sequence which does not contribute the cell adhesion inhibition activity of Hep 2 shows undesirable biological activity. Also it is very complicated procedure to remove urea after being treated with urea and denaturated. Furthermore, high molecular weight molecules can be easily digested in vivo. Due to the above reasons, it is difficult that Hep 2 denaturated with urea it self is used as a medicine.

The peptide having cell adhesion inhibition activity of the present invention is characterized in that it includes the sequence described as the sequence ID No.13 of the sequence listing, and the sequence has a sequence which is absolutely different from the reported peptide sequences having RGD sequence (Humphries, M. J. et al., Science, 233, 467, 1986, Komazawa, H. et al., Biol. Pharm. Bull., 16, 10, 997, 1993). Thus, the cell adhesion inhibition activity of the peptide of the present invention seems not to be mediated by the receptor which is effected by the RGD peptide (Pytela, R. et al., Cell, 48, 549, 1987). Therefore, the peptide of the present invention is a novel peptide sequence having cell adhesion inhibition activity. Furthermore, the cell adhesion inhibition activity is confirmed to be several times stronger than that of the RGD peptide (see the following Examples).

The amino acids sequence described as the sequence ID No. 13 of the sequence listing is a common sequence among the peptides which shows the cell adhesion inhibition activity found by the inventors. Therefore, the peptide having cell adhesion inhibition activity of the present invention is characterized in that it includes the amino acids sequence described as the sequence ID No. 13 of the sequence listing. If a peptide has the amino acids sequence described as the sequence ID No. 13 of the sequence listing in an integral times repeated configuration, the peptide is expected to have more stronger cell adhesion inhibition activity. Such a peptide is also covered with the present invention. The peptide of the present invention includes any amino acid sequences as long as it includes the amino acid sequence described as the sequence ID No. 13 of the sequence listing. Such a peptide is also covered with the present invention. The size (the number of amino acids) of the peptide of the present invention is preferably 30 or less amino acids, more preferably 13 - 30 amino acids, in the light of synthesis efficiency, handling, stability and so forth.

One or several amino acids of the peptide of the present invention may be inserted, deleted or substituted as long as it has the cell adhesion inhibition activity according to the present invention. Such a peptide is covered with the present invention. The insertion, deletion or substitution of one or several amino acids can be made by using a conventional methods such as site-directed mutagenesis.

Also, the derivative of the peptide of the present invention which includes the amino acids sequence described as the sequence ID No. 13 of the sequence listing is expected to have the same or similar effect as the above-mentioned cell adhesion inhibitor peptide, and such a peptide (derivative) is also covered with the present invention. As a method for modifying peptide, for example, modification by using polymer such as polyethylene glycol (PEG) or circularization of linear peptide (Saiki, I. et al., Jpn. J. Cancer Res., 84, 558, 1993) can be listed, but it is not limited to those methods. As long as the cell adhesion inhibition activity is not missed, the peptide of the present invention may be modified by modifying a side chain of the amino acid constituting the peptide with utilization of ester bond, ether bond and so forth. Such a peptide is also covered with the present invention.

The peptide of the present invention having the amino acid sequence described as the sequence ID No. 13 of the sequence listing is thought to be highly hydrophobic base on its amino acid sequence. The peptide of the present invention may be modified by hydrophilic amino acid, a hydrophilic group (such as ethylene glycol and polyethylene glycol) and so forth to induce hydrophilic property of the peptide. Such a modified peptide is also covered with the present invention.

The cell adhesion inhibition peptide of the present invention is useful for investigation and care of the various kinds of disease in which cell adhesion participates. As such a disease, cancer, rheumatism, asthma, allergic disease, thrombosis, rejection against transplantation, healing of wound, inflammation, immunological inflammation containing enteritis nephrocalcinosis (such as colitis ulcerosa), and autoimmune disease can be listed, but it is not limited to these diseases.

The cell adhesion inhibition peptide of the present invention can be administrated by intravenous administration, focus administration to an affected part, oral administration and so forth. Also, additives such as stabilizer and solubilizer can be added on administration if needed.

The cell adhesion inhibition peptide of the present invention can be synthesized by, for example, chemical synthesis such as solid phase chemical synthesis, or gene technology using gene recombinant procedure in which a DNA sequence coding amino acid sequence described as the sequence ID No. 13 is inserted into a plasmid vector, micro-organism such as *E. coli* is transformed with the inserted vector and the peptide is produced. Chemical synthesis is generally done by using a commercialized peptide synthesizer. In the synthesis method using gene recombinant technology, for example, a gene coding the cell adhesion inhibition peptide is synthesized by using DNA synthesizer, the

gene is inserted into a known plasmid vector, the resultant recombinant vector is transformed into a host microorganism to form a recombinant and thus the cell adhesion inhibition peptide is produced. Any expression vector for producing a protein can be used as a plasmid vector used in the synthesis method. Further, the host is not limited to microorganism and eucaryotic cells such as COS cell can be used.

The present invention will be described in detail based on the following working examples. However, the scope of the present invention is not limited by these examples.

Experiment 1: Cell adhesion inhibition activity of a synthetic peptide (a system in which cell suspension includes a synthetic peptide)

The peptide (peptide III 14-2) consisting of the amino acid sequence described as the sequence ID No. 8 in the sequence listing was used in Examples 1 - 3.

Hep 2 fragment treated without urea (Hep 2) was used in Comparative Example 1, and Hep 2 fragment treated with urea (Hep 2-urea) was used in Comparative Example 2. Hep 2 fragment was prepared by a method according to the method of Fukai et al. (Biochem. Biophys. Res. Commun., 220, 394-398, 1996). Hep 2 fragment treated without urea was prepared according to the above method and dissolved in phosphate buffer. Urea treated Hep 2 fragment was prepared by treating with urea according to the method of Fukai et al. mentioned above and thereafter dialyzed against phosphate buffer.

The GRGDSP peptide of cell adhesion domain (describes as the sequence ID No. 9 of the sequence listing), which has been known as a cell adhesion inhibition peptide, was used in Comparative Example 3. The GRGDSP peptide (purchased from IWAKI GLASS, Japan) was dissolved in phosphate buffer.

The peptides consisting of the amino acid sequence described as the sequence ID NO. 1 (peptide III 12-1), the sequence ID NO. 2 (peptide III 12-2), the sequence ID No. 3 (peptide III 12-3), the sequence ID NO. 4 (peptide III 13-1), the sequence ID No. 5 (peptide III 13-2), the sequence ID No. 6 (peptide III 13-3) and the sequence ID No. 7 (peptide III 14-1) were used in Comparative Examples 4 - 9, respectively.

All of the peptides which were used in Examples 1 - 3 and Comparative Examples 4 - 9 were synthesized by applying SAWADY TECHNOLOGY CO., LTD., Tokyo, Japan for synthesis and by using a peptide synthesizer (Multiple Peptide Synthesizer (SYRO II), MultiSynTec GmbH). Further, the sequences of the resultant synthesized peptides were confirmed by using a peptide sequencer (Model 476A, Applied Biosystems).

Each of the synthetic peptides used herein is a peptide corresponding to a part of the sequence of Hep 2 fragment which is derived from Heparin binding domain 2 of fibronectin molecule. Each peptide sequence and Hep 2 sequence as well as the relationship between them are showed in Fig. 1.

The above synthetic peptides were used in this experiment after being dissolved in dimethyl sulfoxide (DMSO: Wako Jyunyaku Co.). To adjust a condition of each experiment to the same, DMSO was added into phosphate buffers containing urea-non-treated Hep 2 fragment, urea-treated Hep 2 fragment, GRGDSP peptide and phosphate buffer itself as a control, respectively. The concentration of DMSO of each experiment was adjusted to 0.2 % of final concentration. Inhibition activity against cell adhesion was evaluated according to the method of Fukai et al. (Biochem. Biophys. Res. Commun., 220, 394-398, 1996). In other words, the inhibition activity against cell adhesion of A375 melanoma cell to a plate was measured under the condition with or without the above various synthetic peptides. A375 melanoma cell (American Type Culture Collection: ATCC) was suspended in Dulbecco's modified eagle medium (DMEM: Gibco Co.) including 0.1 % Ovalbumin (Wako Jyunyaku Co.) to be a concentration of 2×10^5 cells/ml. Before preparing cell suspension, a plate was coated with fibronectin according to the method of Miekka et al. Specifically, phosphate buffer containing 10 µg/ml of fibronectin (FN) was prepared and 100 µl of the resultant phosphate buffer containing fibronectin was added into each well of a 96-well plate and then the plate was incubated at 37 °C, 5 % CO₂ concentration for 60 minutes before being washed with phosphate buffer three times. Fibronectin (FN) was purified according to the method of Miekka et al. (Thromb. Res., 27, 1-14, 1982). One hundred (100) µl of the above cell suspension was added into each well of the plate and the plate was incubated at 37 °C, 5 % CO₂ concentration for 60 minutes (Control was without any synthetic peptide). After fixing cells in each well with 5% of formaldehyde (Wako Jyunyaku Co.), non adhesion cells were removed and calls in five areas selected at random in each well were counted by the use of a microscope.

The addition of Hep 2 fragment and each peptide was carried out by the step in which a solution containing Hep 2 fragment or each peptides dissolved with high concentration was added to the cell suspension to obtain the final concentration of Hep 2 fragment or each peptides shown in Table 1.

The results are shown in Table 1. Table 1 shows that the Hep 2 fragment without urea treatment does not have cell adhesion inhibition activity at all, but after being treated with urea the Hep 2 fragment (Hep 2-urea) has the activity. The peptide III 14-2 has the highest cell adhesion inhibition activity and the activity depends on the concentration of the peptide.

Table 1

Cell adhesion inhibition activity of synthetic peptides		
Fragment or synthetic peptide	Concentration ($\mu\text{g/ml}$)	Cell adhesion inhibition activity*1) (%)
Control	-	0
Comparative Example 1(Hep2)	200	0
Comparative Example 2(Hep2-urea)	50	51
Comparative Example 3(GRGDSP)	200	35
Comparative Example 4(III 12-1)	200	11
Comparative Example 5(III 12-2)	200	13
Comparative Example 6(III 12-3)	200	7
Comparative Example 7(III 13-1)	200	4
Comparative Example 8(III 13-2)	200	11
Comparative Example 9(III 13-3)	200	9
Comparative Example 10(III 14-1)	200	11
Example 1 (III 14-2)	200	71
Example 2 (III 14-2)	20	44
Example 3 (III 14-2)	2	20

*1) The cell adhesion inhibition activity is calculated by counting the number of adhesion cells of the control as 100%, obtaining a percentage of a number of adhesion cells of each of the examples and the comparative examples in which each fragment or peptide is used in comparison with the number of the control (100%), and subtracting the percentage thus obtained from 100.

Experiment 2: Cell adhesion inhibition activity of a synthetic peptide (a system in which a plate is coated with fibronectin solution containing a synthetic peptide)

It is evaluated that which part of the sequence of the peptide III 14-2 contributes its cell adhesion inhibition activity.

The peptides consisting of the amino acid sequence described as the sequence ID NO. 8 (peptide III 14-2), the sequence ID NO. 10 (peptide III 14-2A), the sequence ID No. 11 (peptide III 14-2B) and the sequence ID No. 12 (peptide III 14-2C) were used as the synthetic peptide.

The peptide III 12-2 used in Experiment 1 was used as a control.

All of the peptides which were used in this experiment were synthesized by applying SAWADY TECHNOLOGY CO., LTD., Tokyo, Japan for synthesis and by using a peptide synthesizer (Multiple Peptide Synthesizer (SYRO II), MultiSynTec GmbH). Further, the sequences of the resultant synthesized peptides were confirmed by using a peptide sequencer (Model 476A, Applied Biosystems).

Each of the synthetic peptides used herein was the peptide corresponding to a part of the sequence of Hep 2 fragment which was derived from the heparin binding domain 2 of fibronectin molecule. The relationship among sequences of the peptides is shown in Fig. 2.

All of the synthetic peptides used herein were peptides modified with maleimide-activated Keyhole Limpet Hemocyanine (KLH: PIERCE Co., USA) in order to improve their solubility into phosphate buffer and to coat them to a plate completely (this modification does not influence cell adhesion inhibition activity at all).

The above various KLH modified synthetic peptides were added to a solution containing fibronectin to obtain a given concentration, then the procedure of Experiment 1 were repeated with the exception that the resultant solution were added into 96-well plate to coat it. The experiment without any synthetic peptide is a control (the concentration of a peptide was 0 $\mu\text{g/ml}$). The results are shown in Fig. 3. Fig. 3 shows that, among three peptides containing a part of the peptide III 14-2 (peptides III 14-2A, III 14-2B and III 14-2C), the peptide III 14-2A has very little cell adhesion inhibition activity, while the other two peptides (peptides III 14-2B and III 14-2C) have the same strong cell adhesion inhibition activity as that of the peptide III 14-2. These results indicate that the common amino acid sequence among these three

peptides, peptide III 14-2, peptide III 14-2B and peptide 14-2C (amino acid sequence of YTIYVIAL described as the sequence ID No. 13 of the sequence listing) has cell adhesion inhibition activity.

5 The peptide of the present invention has cell adhesion inhibition activity and, therefore, the peptide can be applied to investigation and therapy of various kind of disease in which cell adhesion participates in. Further, because the peptide of the present invention has cell adhesion inhibition activity the same as or more stronger than that of known cell adhesion inhibition materials and has short length, the synthesis and handling of the peptide of the present invention are easy. Furthermore, the peptide of the present invention is useful as an agent inhibiting cancer metastasis because the peptide has inhibition activity against cancerous cell adhesion.

10 From the invention thus described, it will be obvious that the invention may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications as would be obvious to one skilled in the art are intended for inclusion within the scope of the following claims.

The basic Japanese Application No.8-248247 (248247/1996) filed on September 19, 1996 is hereby incorporated by reference.

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

Sequence ID No.: 1

Sequence length: 31

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1600 - 1630) of Heparin
binding site of human plasma fibronectin

Sequence description:

1 5 10
Ala Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr
15 20 25
Pro Thr Ser Leu Ser Ala Gln Trp Thr Pro Pro Asn Val Gln
25 30
Leu Thr Gly

Sequence ID No.: 2

Sequence length: 22

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1647 - 1668) of Heparin
binding site of human plasma fibronectin

Sequence description:

1 5 10
Ile Asn Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly
15 20
Leu Met Val Ala Thr Lys Trp Glu

Sequence ID No.: 3
5 Sequence length: 23
Sequence type: amino acid
Molecular type: synthetic peptide
10 Feature: Sequence of a part (1669 - 1691) of Heparin
binding site of human plasma fibronectin
15 Sequence description:
1 5 10
Val Ser Val Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg Pro
20 15 20
Ala Gln Gly Val Val Thr Thr Leu Glu

25 Sequence ID No.: 4
Sequence length: 29
30 Sequence type: amino acid
Molecular type: synthetic peptide
35 Feature: Sequence of a part (1692 - 1720) of Heparin
binding site of human plasma fibronectin
Sequence description:
40 1 5 10
Asn Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp Ala Thr
15 20 25
45 Glu Thr Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr
50 Ile

55

Sequence ID No.: 5

Sequence length: 25

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1721 - 1745) of Heparin
binding site of human plasma fibronectin

Sequence description:

1	5	10
Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr		
15	20	25
Pro Ile Gln Arg Thr Ile Lys Pro Asp Val Arg		

Sequence ID No.: 6

Sequence length: 25

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1746 - 1770) of Heparin
binding site of human plasma fibronectin

Sequence description:

1	5	10
Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp Tyr Lys		
15	20	25
Ile Tyr Leu Tyr Thr Leu Asn Asp Asn Ala Arg		

Sequence ID No.: 7

Sequence length: 29

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1787 - 1815) of Heparin
binding site of human plasma fibronectin

Sequence description:

1 5 10

Asn Leu Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu Val

15 20 25

Ser Trp Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile

Lys

Sequence ID No.: 8

Sequence length: 21

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1835 - 1855) of Heparin
binding site of human plasma fibronectin

Sequence description:

1 5 10

Thr Glu Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu Tyr

15 20

Thr Ile Tyr Val Ile Ala Leu

Sequence ID No.: 9

Sequence length: 6

Sequence type: amino acid

5 Molecular type: synthetic peptide

Feature: Sequence of cell binding site of human plasma
fibronectin

10 Sequence description:

1 5

15 Gly Arg Glu Asp Ser Pro

Sequence ID No.: 10

20 Sequence length: 13

Sequence type: amino acid

25 Molecular type: synthetic peptide

Feature: Sequence of a part (1829 - 1841) of Heparin
binding site of human plasma fibronectin

30 Sequence description:

1 5 10

35 Arg Pro Arg Pro Gly Val Thr Glu Ala Thr Ile Thr Gly

Sequence ID No.: 11

40 Sequence length: 19

Sequence type: amino acid

45 Molecular type: synthetic peptide

Feature: Sequence of a part (1842 - 1860) of Heparin
binding site of human plasma fibronectin

50 Sequence description:

1 5 10

55

Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu

15

Lys Asn Asn Gln Lys

Sequence ID No.: 12

Sequence length: 13

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1848 - 1860) of Heparin

binding site of human plasma fibronectin

Sequence description:

1 5 10

Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn Asn Gln Lys

Sequence ID No.: 13

Sequence length: 8

Sequence type: amino acid

Molecular type: synthetic peptide

Feature: Sequence of a part (1848 - 1855) of Heparin

binding site of human plasma fibronectin

Sequence description:

1 5

Tyr Thr Ile Tyr Val Ile Ala Leu

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Hisamitsu Pharmaceutical Co., Inc.
- (B) STREET: 408, Tashirodaikan-machi
- (C) CITY: Tosu-shi
- (D) STATE: Saga
- (E) COUNTRY: Japan
- (F) POSTAL CODE (ZIP): 841

(ii) TITLE OF INVENTION: BIOLOGICALLY ACTIVE PEPTIDE AND CANCER METASTASIS INHIBITOR

(iii) NUMBER OF SEQUENCES: 13

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 97116293.8

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ala Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr Pro Thr
1 5 10 15

Ser Leu Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu Thr Gly
20 25 30

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ile Asn Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly Leu Met
1 5 10 15

Val Ala Thr Lys Trp Glu
20

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Val Ser Val Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg Pro Ala Gln
1 5 10 15

Gly Val Val Thr Thr Leu Glu
20

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Asn Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp Ala Thr Glu Thr
1 5 10 15

Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr Ile
20 25

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro Ile
1 5 10 15

Gln Arg Thr Ile Lys Pro Asp Val Arg
20 25

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp Tyr Lys Ile Tyr
1 5 10 15

Leu Tyr Thr Leu Asn Asp Asn Ala Arg
20 25

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Asn Leu Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu Val Ser Trp
1 5 10 15

Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys
20 25

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Thr Glu Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu Tyr Thr Ile
 1 5 10 15
 Tyr Val Ile Ala Leu
 20

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Gly Arg Glu Asp Ser Pro
 1 5

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Arg Pro Arg Pro Gly Val Thr Glu Ala Thr Ile Thr Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn
1 5 10 15

Asn Gln Lys

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn Asn Gln Lys
1 5 10

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Tyr Thr Ile Tyr Val Ile Ala Leu
1 5

Claims

1. A biologically active peptide having a number of amino acids of 30 or less, comprising an amino acid sequence described as sequence ID No. 13 in the sequence listing and having cell adhesion inhibition activity.
2. A biologically active peptide according to claim 1, wherein the number of amino acids is at least 13.
3. A biologically active peptide having a number of amino acids of 30 or less, comprising an amino acid sequence described as sequence ID No. 8 in the sequence listing and having cell adhesion inhibition activity.
4. A biologically active peptide having a number of amino acids of 30 or less, comprising an amino acid sequence

described as sequence ID No. 11 in the sequence listing and having cell adhesion inhibition activity.

5. A biologically active peptide having a number of amino acids of 30 or less, comprising an amino acid sequence described as sequence ID No. 12 in the sequence listing and having cell adhesion inhibition activity.

5

6. A biological active peptide derived from the peptide according to any one of claims 1 - 5 by insertion, deletion or substitution of one or several amino acids and having the cell adhesion inhibition activity.

10

7. A cancer metastasis inhibitor comprising a biological active peptide according to any one of claims 1 - 5.

15

20

25

30

35

40

45

50

55


Fig.1

Fig.2






Fig.3



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 0 837 074 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
16.09.1998 Bulletin 1998/38

(51) Int. Cl.⁶: C07K 14/78

(43) Date of publication A2:
22.04.1998 Bulletin 1998/17

(21) Application number: 97116293.8

(22) Date of filing: 18.09.1997

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV RO SI

(72) Inventors:
• Fukai, Fumio
Tokorozawa-shi, Saitama 359 (JP)
• Katayama, Takashi
Tokyo 169 (JP)

(30) Priority: 19.09.1996 JP 248247/96

(71) Applicant:
HISAMITSU PHARMACEUTICAL CO., INC.
Tosu-shi Saga-ken (JP)


(74) Representative:
Grünecker, Kinkeldey,
Stockmair & Schwanhäusser
Anwaltssozietät
Maximilianstrasse 58
80538 München (DE)

(54) Biologically active fibronectin fragment as cancer metastasis inhibitor

(57) The biologically active peptide of the invention has a number of amino acids of 30 or less, comprises an amino acid sequence described as sequence ID No. 13 (Tyr Thr Ile Tyr Val Ile Ala Leu) in the sequence listing

and has cell adhesion inhibition activity. The peptide of the invention has short length and, therefore, synthesis and handling of the peptide are easy.

Fig.2



EP 0 837 074 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 97 11 6293 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P,X	F. FUKAI ET AL: "Identification of the anti-adhesive site buried within the heparin-binding domain of fibronectin" JOURNAL OF BIOCHEMISTRY, vol. 121, no. 2, February 1997, TOKYO JP, pages 189-192, XP002051467 * the whole document *	1-7	C07K14/78
X	M.D. PIERSCHBACHER ET AL: "Cell attachment activity of fibronectin can be duplicated by small synthetic fragment of the molecule" NATURE, vol. 309, no. 3, 1984, pages 30-33, XP002051568 * the whole document *	6	
D,A	F. FUKAI: "Fibronectin harbors anticell adhesive activity" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS., vol. 220, 1996, ORLANDO, FL US, pages 394-398, XP002051468 --- -/--		TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search: see sheet C</p>			
Place of search THE HAGUE		Date of completion of the search 12 January 1998	Examiner Cervigni, S
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 62 (P04C07)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 97 11 6293

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	US 5 019 646 A (FURCHT LEO T ET AL) ---		
A	WO 94 13692 A (UNIV MINNESOTA ;US OF AMERICA AS REPRESENTED (US)) * figure 11 * -----		
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 97 11 6293

Claim(s) searched completely:
1-5, 7

Claim(s) searched incompletely:
6

Reason for the limitation of the search:

The scope of claim 6 is unclear and speculative. The claim lacks any structurally constant entity or even any indication concerning the (minimal) size of the peptide, e.g. it even includes dipeptides. Furthermore, the available experimental data lack any sequence exemplifying said claimed compounds. Therefore this claim can also not be considered to represent a permissible generalisation which is fairly based on experimental evidence, that is, it is also not adequately supported by the description. (Art. 84 EPC)

In conclusion, a meaningful and economically feasible search, although directed to compounds having peptidic structure and exhibiting cell adhesion inhibiting activity, could not encompass the complete subject-matter of claim 6 (Rule 45 EPC).